

Hilbert C^* -modules, Lecture 3

①

Chapter 4: Tensor Products

• Throughout, we consider only spatial C^* -algebra tensor products

• $A \otimes B$ will denote the algebraic tensor product.

Here are some results that may or may not be useful for these notes: For C^* -algebras A & B , we have

(i) $A^* \otimes B^*$ embeds in $(A \otimes B)^*$

(ii) For $r \in A \otimes B$, the norm of r is

$$\|r\|^2 = \sup \left\{ \frac{(p \otimes \sigma)(t^* r^* r t)}{(p \otimes \sigma)(t^* t)} : p \in S(A), \sigma \in S(B) \right.$$

where $S(A)$ & $S(B)$ are the set of states $t \in A \otimes B, (p \otimes \sigma)(t^* t) \neq 0$ $\left. \right\}$

(iii) If π & π' are faithful reps of A, B on $\mathcal{H}$ and $\mathcal{K}$, then $\pi \otimes \pi'$ are faithful reps of $A \otimes B$ on $\mathcal{H} \otimes \mathcal{K}$

(2)

(iv) The GNS representation of ρ and σ , for states ρ and σ on A and B respectively, are unitarily equivalent to the representation $\Pi_\rho \otimes \Pi_\sigma$, where Π_ρ, Π_σ are the GNS rep of ρ and σ respectively.

Lemma 4.1 Let E be a Hilbert A -module & t be a bounded A -linear operator on E . TFAE

(i) t is a positive

(ii) $\langle x, tx \rangle \geq 0 \quad \forall x \in E$

Proof: Recall the polarisation identity

$\langle x, y \rangle = \frac{1}{4} \sum_{n=0}^3 i^n \langle x + (i^n)y, x + (-i^n)y \rangle$. So, to show t is self-adjoint, it suffices to show that $\forall x \in E$, $\langle x, tx \rangle = \langle tx, x \rangle$. If $t \geq 0$ in $\mathcal{L}(E)$, then $\langle t^{1/2}x, t^{1/2}x \rangle = \|t^{1/2}x\|^2 \geq 0$ in A . Conversely, by the comment above, if $\langle x, tx \rangle \geq 0 \quad \forall x \in E$, then t is self-adjoint, because $\langle x, tx \rangle^* = \langle tx, x \rangle = \langle x, tx \rangle$. Write $t = r - s$ where $r, s \geq 0$ & $rs = 0$. For $x \in E$, we have $\langle sx, sx \rangle \geq 0$ so $\langle x, s^3x \rangle = -\langle x, s^3x \rangle \geq 0$. But $s^3 \geq 0$, so we must have $\langle x, s^3x \rangle = 0$. Therefore $s = 0$ & $t = r \geq 0$. ■

• Recall the notation $E^n = \underbrace{E \oplus E \oplus \dots \oplus E}_n$ and $\mathcal{L}(E^n) \simeq M_n(\mathcal{L}(E))$, $\mathcal{K}(E^n) \simeq M_n(\mathcal{K}(E))$

Lemma 4.2 Let E be a Hilbert A -module. If $x_1, \dots, x_n \in E$ then $X \geq 0$ in $M_n(A)$, where X is the matrix with (i, j) -entries $\langle x_i, x_j \rangle$. Also, if $t \in \mathcal{L}(E)$ & W is the matrix with entries $\langle tx_i, tx_j \rangle$, then $W \leq \|t\|^2 X$.

Proof: Identify $M_n(A)$ with $\mathcal{K}(E^n)$. Then for all $a = (a_1, \dots, a_n)$ in E^n , we have

$$\langle a, Xa \rangle = \sum_{i,j} a_i^* \langle x_i, x_j \rangle a_j = \left\| \sum_i a_i x_i \right\|^2 \geq 0$$

(3)

Lemma 4.2, continued So, by Lemma 4.1, $X \geq 0$. Also,

$$\begin{aligned} \langle a, Wa \rangle &= \sum_{i,j} a_i^* \langle tx_i, tx_j \rangle a_j = \left| \sum_i tx_i a_i \right|^2 \leq \|t\|^2 \sum_i |x_i a_i|^2 \\ &= \|t\|^2 \langle a, Xa \rangle \end{aligned}$$

The inequality follows from prop 1.2.

Thus, $\langle a, (\|t\|^2 X - W)a \rangle \geq 0 \quad \forall a \in A^n$

So again, by lemma 4.1, $W \leq \|t\|^2 X$. ■

Lemma 4.3 Let A, B be C^* -algebras & $n \in \mathbb{N}$. Suppose that the matrix $a = (a_{ij}) \geq 0$ in $M_n(A)$ & $b = (b_{ij}) \geq 0$ in $M_n(B)$. Then

(i) $(a_{ij} \otimes b_{ij}) \geq 0$ in $M_n(A \otimes B)$

(ii) $\sum_{i,j} a_{ij} \otimes b_{ij} \geq 0$ in $A \otimes B$

(iii) If $a \leq c = (c_{ij})$ in $M_n(A)$, and $b \leq d = (d_{ij}) \in M_n(B)$ then

$$(a_{ij} \otimes b_{ij}) \leq (c_{ij} \otimes d_{ij}) \text{ in } M_n(A \otimes B)$$

Proof: (i) Since $a \geq 0$, $a = S^* S$ for $S \in M_n(A)$. Thus $a_{ij} = \sum_k S_{ki}^* S_{kj}$. Similarly $b_{ij} = \sum_k t_{ki}^* t_{kj}$ in B . Thus,

the matrix $(a_{ij} \otimes b_{ij})$ is the sum of matrices ~~$\sum_{m,k} (S_{ki}^* S_{kj} \otimes t_{mi}^* t_{mj})$~~

$\sum_{m,k} (S_{ki}^* S_{kj} \otimes t_{mi}^* t_{mj})$. It suffices to show that each

matrix in the sum is positive. Let $E = A \otimes B$ be the Hilbert $A \otimes B$ module, and $x_i = S_{ki} \otimes t_{mi}$. Consider $x_1, \dots, x_n \in E$ and the matrix $X_{ij} = \langle x_i, x_j \rangle$. Then lemma 4.2 tells us $(X_{ij}) = (S_{ki}^* S_{kj} \otimes t_{mi}^* t_{mj})$ is a positive matrix, so we are done. ■

(4)

Lemma 4.3, continued

(ii) Let (u_α) and (v_β) be approximate units for A, B respectively. Also, let

$$e_{\alpha\beta} = (\underbrace{u_\alpha \otimes v_\beta, u_\alpha \otimes v_\beta, \dots, u_\alpha \otimes v_\beta}_{n \text{ times}}) \in (A \otimes B)^n$$

It follows that from (i) that $\langle e_{\alpha\beta}, (a_{ij} \otimes b_{ij}) e_{\alpha\beta} \rangle \geq 0$
take the limit to get (ii).

(iii) By (i), the matrices $((c_{ij} - a_{ij}) \otimes b_{ij})$ and $(\bar{c}_{ij} \otimes (d_{ij} - b_{ij}))$ are both positive in $M_n(A \otimes B)$, hence so is their sum.

Constructing Exterior tensor product of Hilbert C^* -modules

Throughout, A & B are C^* -algebras, and E is a Hilbert A -module & F a Hilbert B -module.

We define $E \otimes F$ as a Hilbert $(A \otimes B)$ -module. Let $E \odot F$ be the algebraic tensor product as vector spaces. This is a right module over $A \otimes B$ where $x \otimes y (a \otimes b) = xa \otimes yb$.
For $x_1, x_2 \in E$, $y_1, y_2 \in F$, define

$$\langle x_1 \otimes y_1, x_2 \otimes y_2 \rangle = \langle x_1, x_2 \rangle \otimes \langle y_1, y_2 \rangle$$

This is a semi-inner-product module over $A \otimes B$. It is positive semi-definite because for $z = \sum x_i \otimes y_i \in E \odot F$, then

$$\langle z, z \rangle = \sum_{i,j} \langle x_i, x_j \rangle \otimes \langle y_i, y_j \rangle$$

By lemma 4.2, the matrices $(\langle x_i, x_j \rangle)$ & $(\langle y_i, y_j \rangle)$ are positive. So by lemma 4.3 (ii), $\langle z, z \rangle \geq 0$.

In fact, it is inner product, but too hard to prove. Will prove in Ch. 6.

Taking the completion of this module as mentioned in chapter 1, gives us the exterior product $E \otimes F$.

Example: H a Hilbert Space, A a C^* -algebra as a Hilbert A -module. Then $H \otimes A$ is a Hilbert $C \otimes A \simeq A$ -module.

Adjointable operators on $E \otimes F$

This section shows how $\mathcal{L}(E) \otimes \mathcal{L}(F)$ embeds in $\mathcal{L}(E \otimes F)$ for Hilbert C^* -modules E and F . Choose $s \in \mathcal{L}(E)$ and $t \in \mathcal{L}(F)$. Define $s \otimes t$ on $E \otimes F$ by

$$(s \otimes t)(x \otimes y) = sx \otimes ty.$$

If $z = \sum_i x_i \otimes y_i \in E \otimes F$, then

$$\begin{aligned} |(s \otimes t)z|^2 &= \sum_{i,j} \langle sx_i \otimes ty_i, sx_j \otimes ty_j \rangle \\ &= \sum_{i,j} \langle sx_i, sx_j \rangle \otimes \langle ty_i, ty_j \rangle \end{aligned}$$

By the 2nd part of lemma 4.2, the matrix inequality $C(\langle sx_i, sx_j \rangle) \leq \|s\|^2 C(\langle x_i, x_j \rangle)$ holds & similarly, $C(\langle ty_i, ty_j \rangle) \leq \|t\|^2 C(\langle y_i, y_j \rangle)$. Then, by lemma 4.3 (iii), we have

$$\|s\|^2 \|t\|^2 (C(\langle x_i, x_j \rangle) \otimes C(\langle y_i, y_j \rangle)) - C(\langle sx_i, sx_j \rangle \otimes \langle ty_i, ty_j \rangle) \geq 0$$

using the proof of lemma 4.3, we can turn this into a sum, to get

$$|(s \otimes t)z|^2 = \sum_{i,j} \langle sx_i, sx_j \rangle \otimes \langle ty_i, ty_j \rangle \leq \|s\|^2 \|t\|^2 \sum_{i,j} \langle x_i, x_j \rangle \otimes \langle y_i, y_j \rangle$$

Thus $\|(s \otimes t)z\|^2 \leq \|s\|^2 \|t\|^2 \|z\|^2$. This implies $\|(s \otimes t)z\| \leq \|s\| \|t\| \|z\|$. Therefore, we can extend $s \otimes t$ to be a bounded operator on all of $E \otimes F$.

It is not hard to show that in fact the extension belongs to $\mathcal{L}(E \otimes F)$, with its adjoint being the extension of $S^* \otimes t^*$. Therefore, we have a $*$ -homomorphism $j: \mathcal{L}(E) \otimes \mathcal{L}(F) \rightarrow \mathcal{L}(E \otimes F)$. With a little more work which we omit (see pg 36, 37), one can show j is bounded, and so extends to a $*$ -homomorphism $j: \mathcal{L}(E) \otimes \mathcal{L}(F) \rightarrow \mathcal{L}(E \otimes F)$.

In fact, j is injective. This gives a canonical embedding of $M(A) \otimes M(B)$ into $M(A \otimes B)$ by treating them as Hilbert C^* -modules. j is only surjective if one of the modules is finite dimensional.

Example: Let H be an infinite-dimensional Hilbert space. By the above, $BCH \otimes BCH$ embeds into $BCH \otimes H$, yet, the flip map $\mathcal{L}(g \otimes \eta) = \eta \otimes g$ is not in $BCH \otimes BCH$.

Interior tensor product

Let E be a Hilbert A -module, F a Hilbert B -module. Suppose also we have a $*$ -homomorphism $\phi: A \rightarrow \mathcal{L}(F)$. We want to construct $E \otimes_{\phi} F$ as a Hilbert B -module. To do this, we need to introduce some more concepts.

Lemma 4.4 Suppose E a Hilbert A -module, $x \in E$, and $0 < \alpha < 1$. Then there exists $w \in E$ s.t. $x = w |x|^{\alpha}$.

Proof: For any continuous function f on the spectrum of $|x|$, we have


$$\begin{aligned} \|x f(|x|)\| &= \|f(|x|)^* \langle x, x \rangle f(|x|)\|^{1/2} = \| |x| f(|x|) \| \\ &= \sup \{ |\lambda f(\lambda)| : \lambda \in \text{sp}(|x|) \} \end{aligned}$$

For $n \geq 1$, define a function g_n by

$$g_n(\lambda) = \begin{cases} n^{\alpha} & \text{if } \lambda \leq 1/n \\ \lambda^{-\alpha} & \text{if } \lambda > 1/n \end{cases}$$

(7)

Lemma 4.4, cont. so the graph of g_n looks like



note that $(\lambda g_n(|X|))$ is a $\frac{1}{n}$ Cauchy sequence in E because

$$\sup_{\lambda \in \text{sp}(|X|)} |\lambda(g_n(\lambda) - g_m(\lambda))| \leq \max(n^{\alpha-1}, m^{\alpha-1})$$

and $\alpha < 1$, so this goes to 0. Thus, $\exists$ limit w in E . Adjoining 1 to A if needed, & since $1x = 1x = x$ in E , we have

$$\|x g_n(|X|) |X|^\alpha - x\| = \|x(g_n(|X|)(|X|^\alpha - 1))\|$$

since $\sup_{\lambda \in \text{sp}(|X|)} \{ |\lambda(g_n(\lambda)|X|^\alpha - 1)| : \lambda \in \text{sp}(|X|) \} \rightarrow 0$

as $n \rightarrow \infty$, we have $w|X|^\alpha = x$. ■

Definition: If $\varphi: A \rightarrow B$ a linear map between C^* -algebras, $\varphi^{(n)}$ is a map from $M_n(A)$ to $M_n(B)$ obtained by $\varphi^{(n)}([a_{ij}]) = [\varphi(a_{ij})]$. φ is said to be completely positive if $\varphi^{(n)}$ positive map for all $n \in \mathbb{N}$.

If φ $*$ -homomorphism, then it is completely positive.

Let E be an A -Hilbert module. E^n can be made into an $M_n(A)$ -module as follows: for $a \in M_n(A)$, $x = (x_1, \dots, x_n) \in E^n$ let $xa = (x_1, \dots, x_n)a$ where multiplication is how you would normally multiply a $1 \times n$ matrix by an $n \times n$ matrix. So you consider elements of E^n as 'row vectors'. The inner product $\langle x, y \rangle$ is the matrix $(\langle x_i, y_j \rangle)$.

(8)

The fact that $\langle x, x \rangle \geq 0$ in this case follows from Lemma 4.2.

Constructing the Interior tensor product

Let E, f and $\phi: A \rightarrow \mathcal{L}(f)$ as before. f can be seen as a left A -module, with action given by $(a, y) \mapsto \phi(a)y$.

Take the algebraic vector space tensor product $E \otimes f$, and quotient out by the subspace generated by

$$x a \otimes y - x \otimes \phi(a)y \quad (x \in E, y \in f, a \in A) \quad (4.3)$$

B acts on this space by $(x \otimes y)b = x \otimes (yb)$. We denote this B -module by $E \otimes_A f$.

Proposition 4.5 With A, B, E, f & ϕ as above, $E \otimes_A f$ is an inner-product B -module with inner product

$$\langle x_1 \otimes y_1, x_2 \otimes y_2 \rangle = \langle y_1, \phi(\langle x_1, x_2 \rangle) y_2 \rangle \quad (4.4)$$

Proof: If $z = \sum_{i=1}^n x_i \otimes y_i \in E \otimes f$, then

$$\begin{aligned} \langle z, z \rangle &= \sum_{i,j} \langle y_i, \phi(\langle x_i, x_j \rangle) y_j \rangle \\ &= \langle y, \phi^{(n)}(X) y \rangle \end{aligned}$$

where $y = (y_1, \dots, y_n) \in f^n$, and $X \in M_n(A)$, where $X_{ij} = \langle x_i, x_j \rangle$. By Lemma 4.2, $X \geq 0$. Since ϕ is a completely positive map, $\phi^{(n)}(X) \geq 0$, so $\langle z, z \rangle \geq 0$.

Thus, (4.4) turns $E \otimes f$ into a semi inner-product B module. Let

$$N = \{ z \in E \otimes f : \langle z, z \rangle = 0 \}$$

Proposition 4.3 continued. We will show N is the same as the subspace generated by (4.3), so the semi inner-product factors through as an inner-product on $E \otimes_A F$.

Is $z = x \otimes y - x \otimes \phi(a)y$, then

$$\begin{aligned} \langle z, z \rangle &= \langle x \otimes y, x \otimes y \rangle - \langle x \otimes y, x \otimes \phi(a)y \rangle - \\ &\quad - \langle x \otimes \phi(a)y, x \otimes y \rangle + \langle x \otimes \phi(a)y, x \otimes \phi(a)y \rangle \\ &= \langle y, \phi(a^* |x|^2 a) y \rangle - \langle y, \phi(a^* |x|^2 a) y \rangle - \langle y, \phi(a^* |x|^2 a) y \rangle \\ &\quad + \langle y, \phi(a^* |x|^2 a) y \rangle = 0 \end{aligned}$$

Conversely, suppose $z = \sum x_i \otimes y_i \in N$. Then using previous notation,

$$\langle z, z \rangle = \langle y, \phi^{(n)}(X) y \rangle = 0$$

for $y \in F^n$, $X \in M_n(A)$. Let $T = \phi^{(n)}(X)$. Then $T \geq 0$ in $M_n(\mathcal{L}(F)) \cong \mathcal{L}(F^n)$ and $T^{1/4} y = 0$.

Treat E^n as a $M_n(A)$ -module as mentioned earlier. let $\underline{x} = (x_1, \dots, x_n)$ then little $|\underline{x}| = \|\underline{x}\|^{1/2}$ (big $X_{ij} = \langle \underline{x}_i, \underline{x}_j \rangle$). By lemma 4.4, $\exists \omega \in E^n$ s.t. $\omega |\underline{x}|^{1/2} = \omega X^{1/4} = \underline{x}$. write c_{ij} for matrix elements of $X^{1/4}$. Then $T^{1/4} = \phi^{(n)}(X^{1/4})$, so $(T^{1/4})_{ij} = \phi(c_{ij})$. So, we have

$$T^{1/4} y = 0 \Rightarrow \forall j=1, \dots, n, \sum_i \phi(c_{ij}) y_j = 0$$

and,

$$\underline{x}_j = \sum_i \omega_i c_{ij}, \text{ since } \underline{\omega} X^{1/4} = \underline{x}.$$

So, $z = \sum_j x_j \otimes y_j$, using the formulae above, we can write

$$z = \sum_{i,j} \omega_i c_{ij} \otimes y_j - \omega_i \otimes \phi(c_{ij}) y_j, \text{ so } z \text{ is of the form of (4.3). } \blacksquare$$

(10)

Completion of $E \otimes_A F$ with this inner product is the Hilbert B -module denoted $E \otimes_\phi F$. We denote elements of $E \otimes_\phi F$ by $x \otimes y$.

Example: $E = H \otimes A$, for a Hilbert space H , a C^* -algebra A . ~~Let~~ Write $H_A = H \otimes A$. Let F be a Hilbert B -module & suppose $\phi \in \text{Mor}(A, K(E))$.

Then linear combinations of elements of the form $(g \otimes a) \otimes y$ are dense in $H_A \otimes_\phi F$. By nondegeneracy of ϕ , sum of the form $g \otimes \phi(a)y$ are dense in $H \otimes F$. One can show that

$$\| \sum_i (g_i \otimes a_i) \otimes y_i \| = \| \sum_i g_i \otimes \phi(a_i) y_i \| \quad (4.5)$$

So the map $(g \otimes a) \otimes y \mapsto g \otimes \phi(a)y$ extends to a unitary map from $H_A \otimes_\phi F$ to $H \otimes F$. So the two modules are unitarily equivalent.

Using this example when $F=B$, we get $H_A \otimes_\phi B \simeq H_B$.

Now suppose E is a Hilbert A -module, $t \in \mathcal{L}(E)$, F a Hilbert B -module & $\phi: A \rightarrow \mathcal{L}(F)$ $*$ -hom. The map defined by $x \otimes y \mapsto tx \otimes y$ extends to a bounded map on $E \otimes_\phi F$. This map is denoted by $\phi_*(t) \in \mathcal{L}(F \otimes_\phi F)$.

The map $\phi_*: \mathcal{L}(E) \rightarrow \mathcal{L}(F \otimes_\phi F)$ is a unital nondegenerate $*$ -hom. ϕ_* is injective iff ϕ is injective.